

[preprint] aastex $\text{H}\alpha$ $\text{H}\beta$ HeI $\lambda 6678$ $[\text{O} \text{I}]$ $\lambda 5007$ $[\text{O} \text{I}]$ $\lambda 4959$ $[\text{O} \text{I}]$ $\lambda 4959+5007$ $[\text{N} \text{I}]$ $\lambda 6584$ $[\text{N} \text{I}]$ $\lambda 6548+6584$ $[\text{S} \text{I}]$ $\lambda 6717+6731$ $[\text{S} \text{I}]$ $\lambda 6717$ $[\text{S} \text{I}]$ $\lambda 6731$ $W(\text{H}\alpha)$ $W(\text{H}\beta)$ $W([\text{O} \text{I}] \lambda 5007)$ $W(\text{WR } \lambda 4686)$ $L(\text{WR } \lambda 4686)$ $L(\text{H}\beta)$
 To appear in the Astronomical Journal MacKenty et al. 2000 HST/WFPC2 and VLA Observations of the Ionized Gas in NGC 4214 document *et al.*

HST/WFPC2 and VLA Observations of the Ionized Gas in the Dwarf Starburst Galaxy NGC 4214
 The Ionized Gas in NGC 4214

John W. MacKenty², Jesús Maíz-Apellániz^{2,3}, Christopher E. Pickens⁴, Colin A. Norman^{2,4}, Nolan R. Walborn²

¹Based on observations with the NASA/ESA Hubble Space Telescope and the NRAO Very Large Array. The HST observations were obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract No. NAS5-26555. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

²Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, U.S.A.

³Laboratorio de Astrofísica Espacial y Física Fundamental-INTA, Apdo. Postal 50727, E-28080 Madrid, Spain.

⁴Henry A. Rowland Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, Maryland 21218, U.S.A.

abstract

We present new and narrow band images of the starbursting dwarf galaxy NGC 4214, obtained with the Wide Field and Planetary Camera (WFPC2) onboard the Hubble Space Telescope (HST), together with VLA observations of the same galaxy. The HST images resolve features down to physical scales of 2 – 5 pc, revealing several young (< 10 Myr) star forming complexes of various ionized gas morphologies (compact knots, complete or fragmentary shells) and sizes ($\sim 10 - 200$ pc). Our results are consistent with a uniform set of evolutionary trends: The youngest, smaller, filled regions that presumably are those just emerging from dense star forming clouds, tend to be of high excitation and are highly obscured. Evolved, larger shell-like regions have lower excitation and are less extincted due of the action of stellar winds and supernovae. In at least one case we find evidence for induced star formation which has led to a two-stage starburst. Age estimates based on measurements do not agree with those inferred from wind-driven shell models of expanding H₂ regions. The most likely explanation for this effect is the existence of an ≈ 2 Myr delay in the formation of superbubbles caused by the pressure exerted by the high density medium in which massive stars are born. We report the detection of a supernova remnant embedded in one of the two large H₂ complexes of NGC 4214. The dust in NGC 4214 is not located in a foreground screen but is physically associated with the warm ionized gas.





